

**COEFFICIENT ESTIMATE OF MA-MINDA TYPE
BI-BAZILEVIČ FUNCTIONS OF COMPLEX ORDER
INVOLVING SRIVASTAVA-ATTIYA OPERATOR**

G. MURUGUSUNDARAMOORTHY, C. SELVARAJ, O. S. BABU

ABSTRACT. In this paper, we introduce and investigate a new subclass of the function class Σ of bi-univalent functions defined in the open unit disk, which are associated with the Hurwitz-Lerch zeta function, satisfying subordinate conditions. Furthermore, we find estimates on the Taylor-Maclaurin coefficients $|a_2|$ and $|a_3|$ for functions in this new subclass. Several (known or new) consequences of the results are also pointed out.

1. INTRODUCTION, DEFINITIONS AND PRELIMINARIES

Let \mathcal{A} denote the class of functions of the form:

$$f(z) = z + \sum_{n=2}^{\infty} a_n z^n, \quad (1)$$

which are analytic in the open unit disk

$$\Delta = \{z : z \in \mathbb{C} \text{ and } |z| < 1\}.$$

Further, by \mathcal{S} we shall denote the class of all functions in \mathcal{A} which are univalent in Δ . Some of the important and well-investigated subclasses of the univalent function class \mathcal{S} include (for example) the class $\mathcal{S}^*(\alpha)$ of starlike functions of order α in Δ and the class $\mathcal{K}(\alpha)$ of convex functions of order α in Δ . It is well known that every function $f \in \mathcal{S}$ has an inverse f^{-1} , defined by

$$f^{-1}(f(z)) = z \quad (z \in \Delta)$$

and

$$f(f^{-1}(w)) = w \quad \left(|w| < r_0(f); r_0(f) \geq \frac{1}{4} \right),$$

where

$$g(w) = f^{-1}(w) = w - a_2 w^2 + (2a_2^2 - a_3)w^3 - (5a_2^3 - 5a_2 a_3 + a_4)w^4 + \cdots \quad (2)$$

2000 *Mathematics Subject Classification*. Primary 30C45.

Key words and phrases. Analytic functions; Univalent functions; Bi-univalent functions; Bi-starlike and bi-convex functions; Bi-Bazilevič functions; Hurwitz-Lerch zeta function; Jung-Kim-Srivastava integral operator; Libera-Bernardi integral operator.

Submitted sep. 15, 2013. Revised Nov. 4, 2013.

A function $f \in \mathcal{A}$ is said to be bi-univalent in Δ if both $f(z)$ and $f^{-1}(z)$ are univalent in Δ . Let Σ denote the class of bi-univalent functions in Δ given by (1).

An analytic function f is subordinate to an analytic function g , written $f(z) \prec g(z)$, provided there is an analytic function w defined on Δ with $w(0) = 0$ and $|w(z)| < 1$ satisfying $f(z) = g(w(z))$. Ma and Minda [18] unified various subclasses of starlike and convex functions for which either of the quantity $\frac{zf'(z)}{f(z)}$ or $1 + \frac{zf''(z)}{f'(z)}$ is subordinate to a more general superordinate function. For this purpose, they considered an analytic function ϕ with positive real part in the unit disk Δ , $\phi(0) = 1$, $\phi'(0) > 0$, and ϕ maps Δ onto a region starlike with respect to 1 and symmetric with respect to the real axis. The class of Ma-Minda starlike functions consists of functions $f \in \mathcal{A}$ satisfying the subordination $\frac{zf'(z)}{f(z)} \prec \phi(z)$. Similarly, the class of Ma-Minda convex functions of functions $f \in \mathcal{A}$ satisfying the subordination $1 + \frac{zf''(z)}{f'(z)} \prec \phi(z)$.

A function f is bi-starlike of Ma-Minda type or bi-convex of Ma-Minda type if both f and f^{-1} are respectively Ma-Minda starlike or convex. These classes are denoted respectively by $\mathcal{S}_{\Sigma}^*(\phi)$ and $\mathcal{K}_{\Sigma}(\phi)$, where $\phi(z)$ is given by

$$\phi(z) = 1 + B_1z + B_2z^2 + B_3z^3 + \dots, \quad (B_1 > 0). \quad (3)$$

The convolution or Hadamard product of two functions $f, h \in \mathcal{A}$ is denoted by $f * h$ and is defined as

$$(f * h)(z) = z + \sum_{n=2}^{\infty} a_n b_n z^n, \quad (4)$$

where $f(z)$ is given by (1) and $h(z) = z + \sum_{n=2}^{\infty} b_n z^n$.

We recall a general Hurwitz-Lerch Zeta function $\Phi(z, s, a)$ defined by (see [25])

$$\Phi(z, s, a) := \sum_{k=0}^{\infty} \frac{z^k}{(k+a)^s} \quad (a \in \mathbb{C} \setminus \{0, -1, -2, \dots\}; s \in \mathbb{C}, \Re(s) > 1 \text{ and } |z| < 1). \quad (5)$$

Several interesting properties and characteristics of the Hurwitz-Lerch zeta function $\Phi(z, s, a)$ can be found in the recent investigations by Choi and Srivastava [5], Ferreira and Lopez [7], Garg et al [10], Lin and Srivastava [16], Lin et al [17] and others.

For the class \mathcal{A} , Srivastava and Attiya [24] (see also Raducanu and Srivastava [22] and Prajapat and Goyal [21]) introduced and investigated linear operator:

$$\mathcal{J}_{\mu, b} : \mathcal{A} \longrightarrow \mathcal{A}$$

defined in terms of the Hadamard product (or convolution) by

$$\mathcal{J}_{\mu, b} f(z) = (\mathcal{G}_{\mu, b} * f)(z) \quad (z \in \Delta; b \in \mathbb{C} \setminus \{0, -1, -2, \dots\}; \mu \in \mathbb{C}; f \in \mathcal{A}), \quad (6)$$

where, for convenience.

$$\mathcal{G}_{\mu, b}(z) = (1+b)^{\mu} [\Phi(z, \mu, b) - b^{-\mu}]. \quad (7)$$

It is easy to observe from (given earlier by [21], [22]) (1), (6) and (7) that

$$\mathcal{J}_{\mu, b} f(z) = z + \sum_{k=2}^{\infty} \Theta_k a_k z^k, \quad (8)$$

where

$$\Theta_k = \left| \left(\frac{1+b}{k+b} \right)^\mu \right| \quad (9)$$

and (throughout this paper unless otherwise mentioned) the parameters μ, b are considered as $\mu \in \mathbb{C}$ and $b \in \mathbb{C} \setminus \{0, -1, -2, \dots\}$.

We note that

- For $\mu = 1$ and $b = \nu (\nu > -1)$ generalized Libera-Bernardi integral operator [23]

$$\begin{aligned} \mathcal{J}_{1,\nu} f(z) &= \frac{1+\nu}{z^\nu} \int_0^z t^{\nu-1} f(t) dt \\ &= z + \sum_{k=2}^{\infty} \left(\frac{\nu+1}{k+\nu} \right) a_k z^k = \mathcal{L}_\nu f(z). \end{aligned}$$

- For $\mu = \sigma (\sigma > 0)$ and $b = 1$, Jung-Kim-Srivastava integral operator [13]

$$\begin{aligned} \mathcal{J}_{\sigma,1} f(z) &= \frac{2^\sigma}{z\Gamma(\sigma)} \int_0^z \left(\log \left(\frac{z}{t} \right) \right)^{\sigma-1} f(t) dt \\ &= z + \sum_{k=2}^{\infty} \left(\frac{z}{k+1} \right)^\sigma a_k z^k = \mathcal{I}_\sigma f(z) \end{aligned}$$

closely related to some multiplier transformations studied by Flett [8].

Recently there has been triggering interest to study bi-univalent function class Σ and obtained non-sharp coefficient estimates on the first two coefficients $|a_2|$ and $|a_3|$ of (1). But the coefficient problem for each of the following Taylor-Maclaurin coefficients:

$$|a_n| \quad (n \in \mathbb{N} \setminus \{1, 2, 3\}; \mathbb{N} := \{1, 2, 3, \dots\})$$

is still an open problem(see[2, 3, 4, 14, 19, 27]). Many researchers (see[9, 11, 15, 26, 28, 29]) have recently introduced and investigated several interesting subclasses of the bi-univalent function class Σ and they have found non-sharp estimates on the first two Taylor-Maclaurin coefficients $|a_2|$ and $|a_3|$.

Several authors have discussed various subfamilies of Bazilevič functions of type λ from various perspective. They discussed it from the perspective of convexity, inclusion theorem, radii of starlikeness and convexity boundary rotational problem, subordination just to mention few. The most amazing thing is that, it is difficult to see any of this authors discussing the coefficient inequalities, and coefficient bounds of these subfamilies of Bazilevič function most especially when the parameter λ is greater than 1 ($\lambda \in \mathbb{R}$). Motivated by the earlier work of Deniz[6] in the present paper we introduce new families of Bazilevič functions of complex order [12] of the function class Σ , involving Hurwitz-Lerch zeta function, and find estimates on the coefficients $|a_2|$ and $|a_3|$ for functions in the new subclasses of function class Σ . Several related classes are also considered, and connection to earlier known results are made.

Definition 1. A function $f \in \Sigma$ given by (1) is said to be in the class $\mathcal{S}_{\Sigma}^{\mu,b}(\gamma, \lambda, \phi)$ if the following conditions are satisfied:

$$1 + \frac{1}{\gamma} \left(\frac{z^{1-\lambda}(\mathcal{J}_{\mu,b}f(z))'}{[\mathcal{J}_{\mu,b}f(z)]^{1-\lambda}} - 1 \right) \prec \phi(z) \quad (10)$$

and

$$1 + \frac{1}{\gamma} \left(\frac{w^{1-\lambda}(\mathcal{J}_{\mu,b}g(w))'}{[\mathcal{J}_{\mu,b}g(w)]^{1-\lambda}} - 1 \right) \prec \phi(w) \quad (11)$$

where $\gamma \in \mathbb{C} \setminus \{0\}$; $\lambda \geq 0$; $z, w \in \Delta$ and the function g is given by(2).

Example 1. If we set $\phi(z) = \frac{1+Az}{1+Bz}$, $-1 \leq B < A \leq 1$, then the class $\mathcal{S}_{\Sigma}^{\mu,b}(\gamma, \lambda, \phi) \equiv \mathcal{S}_{\Sigma}^{\mu,b}(\gamma, \lambda, A, B)$ which is defined as $f \in \Sigma$,

$$1 + \frac{1}{\gamma} \left(\frac{z^{1-\lambda}(\mathcal{J}_{\mu,b}f(z))'}{[\mathcal{J}_{\mu,b}f(z)]^{1-\lambda}} - 1 \right) \prec \frac{1+Az}{1+Bz},$$

and

$$1 + \frac{1}{\gamma} \left(\frac{w^{1-\lambda}(\mathcal{J}_{\mu,b}g(w))'}{[\mathcal{J}_{\mu,b}g(w)]^{1-\lambda}} - 1 \right) \prec \frac{1+Aw}{1+Bw}.$$

Example 2. If we set $\phi(z) = \frac{1+(1-2\alpha)z}{1-z}$, $0 \leq \alpha < 1$ then the class $\mathcal{S}_{\Sigma}^{\mu,b}(\gamma, \lambda, \phi) \equiv \mathcal{S}_{\Sigma}^{\mu,b}(\gamma, \lambda, \alpha)$ which is defined as $f \in \Sigma$,

$$\Re \left[1 + \frac{1}{\gamma} \left(\frac{z^{1-\lambda}(\mathcal{J}_{\mu,b}f(z))'}{[\mathcal{J}_{\mu,b}f(z)]^{1-\lambda}} - 1 \right) \right] > \alpha,$$

and

$$\Re \left[1 + \frac{1}{\gamma} \left(\frac{w^{1-\lambda}(\mathcal{J}_{\mu,b}g(w))'}{[\mathcal{J}_{\mu,b}g(w)]^{1-\lambda}} - 1 \right) \right] > \alpha.$$

On specializing the parameters λ one can state the various new subclasses of Σ as illustrated in the following examples.

Example 3. For $\lambda = 0$ and a function $f \in \Sigma$, given by (1) is said to be in the class $\mathcal{S}_{\Sigma}^{\mu,b}(\gamma, \phi)$ if the following conditions are satisfied:

$$1 + \frac{1}{\gamma} \left(\frac{z(\mathcal{J}_{\mu,b}f(z))'}{\mathcal{J}_{\mu,b}f(z)} - 1 \right) \prec \phi(z) \quad (12)$$

and

$$1 + \frac{1}{\gamma} \left(\frac{w(\mathcal{J}_{\mu,b}g(w))'}{\mathcal{J}_{\mu,b}g(w)} - 1 \right) \prec \phi(w) \quad (13)$$

where $\gamma \in \mathbb{C} \setminus \{0\}$; $z, w \in \Delta$ and the function g is given by(2).

Example 4. For $\lambda = 1$ and a function $f \in \Sigma$, given by (1) is said to be in the class $\mathcal{H}_{\Sigma}^{\mu,b}(\gamma, \phi)$ if the following conditions are satisfied:

$$1 + \frac{1}{\gamma} (\mathcal{J}_{\mu,b}f(z))' - 1 \prec \phi(z) \quad (14)$$

and

$$1 + \frac{1}{\gamma} ((\mathcal{J}_{\mu,b}g(w))' - 1) \prec \phi(w) \quad (15)$$

where $\gamma \in \mathbb{C} \setminus \{0\}$; $z, w \in \Delta$ and the function g is given by(2).

It is of interest to note that for $\gamma = 1$ the class $\mathcal{S}_{\Sigma}^{\mu,b}(\gamma, \lambda, \phi)$ reduces to the following new subclass $\mathcal{B}_{\Sigma}^{\mu,b}(\lambda, \phi)$.

Definition 2. A function $f \in \Sigma$ given by (1) is said to be in the class $\mathcal{B}_{\Sigma}^{\mu,b}(\lambda, \phi)$ if the following conditions are satisfied:

$$\frac{z^{1-\lambda}(\mathcal{J}_{\mu,b}f(z))'}{[\mathcal{J}_{\mu,b}f(z)]^{1-\lambda}} \prec \phi(z) \quad (16)$$

and

$$\frac{w^{1-\lambda}(\mathcal{J}_{\mu,b}g(w))'}{[\mathcal{J}_{\mu,b}g(w)]^{1-\lambda}} \prec \phi(w) \quad (17)$$

where $\lambda \geq 0$; $z, w \in \Delta$ and the function g is given by(2).

For particular values of λ , we have

Example 5. For $\lambda = 0$ and a function $f \in \Sigma$, given by (1) is said to be in the class $\mathcal{B}_{\Sigma}^{\mu,b}(0, \phi) \equiv \mathcal{S}_{\Sigma}^{*,\mu,b}(\phi)$ if the following conditions are satisfied:

$$\frac{z(\mathcal{J}_{\mu,b}f(z))'}{\mathcal{J}_{\mu,b}f(z)} \prec \phi(z) \quad (18)$$

and

$$\frac{w(\mathcal{J}_{\mu,b}g(w))'}{\mathcal{J}_{\mu,b}g(w)} \prec \phi(w) \quad (19)$$

where $z, w \in \Delta$ and the function g is given by(2).

Example 6. For $\lambda = 1$ and a function $f \in \Sigma$, given by (1) is said to be in the class $\mathcal{B}_{\Sigma}^{\mu,b}(1, \phi) \equiv \mathcal{H}_{\Sigma}^{\mu,b}(\phi)$ if the following conditions are satisfied:

$$(\mathcal{J}_{\mu,b}f(z))' \prec \phi(z) \quad (20)$$

and

$$(\mathcal{J}_{\mu,b}g(w))' \prec \phi(w) \quad (21)$$

where $z, w \in \Delta$ and the function g is given by(2).

Remark 3. By setting $\phi(z) = \frac{1+Az}{1+Bz}$, $-1 \leq B < A \leq 1$, (or $\phi(z) = \frac{1+(1-2\alpha)z}{1-z}$, $0 \leq \alpha < 1$) as mentioned in Example 1 and 2 we state some new analogous subclasses $\mathcal{S}_{\Sigma}^{\mu,b}(\gamma, A, B)$; $\mathcal{S}_{\Sigma}^{\mu,b}(\gamma, \alpha)$; $\mathcal{H}_{\Sigma}^{\mu,b}(\gamma, A, B)$; $\mathcal{H}_{\Sigma}^{\mu,b}(\gamma, \alpha)$; $\mathcal{S}_{\Sigma}^{*,\mu,b}(\gamma, A, B)$; $\mathcal{S}_{\Sigma}^{*,\mu,b}(\gamma, \alpha)$; $\mathcal{H}_{\Sigma}^{\mu,b}(\gamma, A, B)$ $\mathcal{H}_{\Sigma}^{\mu,b}(\gamma, \alpha)$ for the classes defined in Examples 3 to 6 respectively .

Remark 4. When $\mathcal{J}_{0,b}f(z) = f(z)$ (that is, $\Theta_k = 1, k = 2, 3, \dots$) the class $\mathcal{S}_{\Sigma}^{\mu,b}(\gamma, \lambda, \phi) \equiv \mathcal{S}_{\Sigma}(\gamma, \lambda, \phi)$, the class of Ma-Minda type bi-Bazilevič functions of complex order; $\mathcal{S}_{\Sigma}^{*,\mu,b}(\gamma, \phi) \equiv \mathcal{S}_{\Sigma}^*(\gamma, \phi)$, the class of Ma-Minda type Bi-starlike functions of complex order; $\mathcal{H}_{\Sigma}^{\mu,b}(\gamma, \phi) \equiv \mathcal{H}_{\Sigma}(\gamma, \phi)$; the class $\mathcal{B}_{\Sigma}^{\mu,b}(\lambda, \phi) \equiv \mathcal{B}_{\Sigma}(\lambda, \phi)$, the class of bi-Bazilevič functions of Ma-Minda type [18] the class $\mathcal{S}_{\Sigma}^{*,\mu,b}(\phi) \equiv \mathcal{S}_{\Sigma}^*(\phi)$, the class of Ma-Minda bi-starlike functions [1] and the class $\mathcal{H}_{\Sigma}^{\mu,b}(\phi) \equiv \mathcal{H}_{\Sigma}(\phi)$ [1, 26].

In the following section we find estimates on the coefficients $|a_2|$ and $|a_3|$ for functions in the above-defined subclasses $\mathcal{S}_{\Sigma}^{\mu,b}(\gamma, \lambda, \phi)$ of the function class Σ .

In order to derive our main results, we shall need the following lemma:

Lemma 5. (see [20]) *If $p \in \mathcal{P}$, then $|p_k| \leq 2$ for each k , where \mathcal{P} is the family of all functions p analytic in Δ for which $\Re(p(z)) > 0$, where $p(z) = 1 + p_1z + p_2z^2 + \dots$ for $z \in \Delta$.*

2. COEFFICIENT BOUNDS FOR THE FUNCTION CLASS $\mathcal{S}_{\Sigma}^{\mu,b}(\gamma, \lambda, \phi)$

We begin by finding the estimates on the coefficients $|a_2|$ and $|a_3|$ for functions in the class $\mathcal{S}_{\Sigma}^{\mu,b}(\gamma, \lambda, \phi)$.

Theorem 6. *Let the function $f(z)$ given by (1) be in the class $\mathcal{S}_{\Sigma}^{\mu,b}(\gamma, \lambda, \phi)$. Then*

$$|a_2| \leq \frac{|\gamma|B_1\sqrt{2B_1}}{\sqrt{|\gamma B_1^2[(\lambda-1)(\lambda+2)\Theta_2^2 + 2(\lambda+2)\Theta_3] - 2(B_2 - B_1)(1+\lambda)^2\Theta_2^2|}} \quad (22)$$

and

$$|a_3| \leq \frac{|\gamma|B_1}{(\lambda+2)\Theta_3} + \left(\frac{|\gamma|B_1}{(1+\lambda)\Theta_2} \right)^2. \quad (23)$$

Proof. Let $f \in \mathcal{S}_{\Sigma}^{\mu,b}(\gamma, \lambda, \phi)$ and $g = f^{-1}$. Then there are analytic functions $u, v : \Delta \rightarrow \Delta$ with $u(0) = 0 = v(0)$, satisfying

$$1 + \frac{1}{\gamma} \left(\frac{z^{1-\lambda}(\mathcal{J}_{\mu,b}f(z))'}{[\mathcal{J}_{\mu,b}f(z)]^{1-\lambda}} - 1 \right) = \phi(u(z)) \quad (24)$$

and

$$1 + \frac{1}{\gamma} \left(\frac{w^{1-\lambda}(\mathcal{J}_{\mu,b}g(w))'}{[\mathcal{J}_{\mu,b}g(w)]^{1-\lambda}} - 1 \right) = \phi(v(w)). \quad (25)$$

Define the functions $p(z)$ and $q(z)$ by

$$p(z) := \frac{1+u(z)}{1-u(z)} = 1 + p_1z + p_2z^2 + \dots$$

and

$$q(z) := \frac{1+v(z)}{1-v(z)} = 1 + q_1z + q_2z^2 + \dots$$

or, equivalently,

$$u(z) := \frac{p(z)-1}{p(z)+1} = \frac{1}{2} \left[p_1z + \left(p_2 - \frac{p_1^2}{2} \right) z^2 + \dots \right] \quad (26)$$

and

$$v(z) := \frac{q(z)-1}{q(z)+1} = \frac{1}{2} \left[q_1z + \left(q_2 - \frac{q_1^2}{2} \right) z^2 + \dots \right]. \quad (27)$$

Then $p(z)$ and $q(z)$ are analytic in Δ with $p(0) = 1 = q(0)$. Since $u, v : \Delta \rightarrow \Delta$, the functions $p(z)$ and $q(z)$ have a positive real part in Δ , and $|p_i| \leq 2$ and $|q_i| \leq 2$.

Using (26) and (27) in (24) and (25) respectively, we have

$$1 + \frac{1}{\gamma} \left(\frac{z^{1-\lambda}(\mathcal{J}_{\mu,b}f(z))'}{[\mathcal{J}_{\mu,b}f(z)]^{1-\lambda}} - 1 \right) = \varphi \left(\frac{1}{2} \left[p_1z + \left(p_2 - \frac{p_1^2}{2} \right) z^2 + \dots \right] \right) \quad (28)$$

and

$$1 + \frac{1}{\gamma} \left(\frac{w^{1-\lambda}(\mathcal{J}_{\mu,b}g(w))'}{[\mathcal{J}_{\mu,b}g(w)]^{1-\lambda}} - 1 \right) = \varphi \left(\frac{1}{2} \left[q_1w + \left(q_2 - \frac{q_1^2}{2} \right) w^2 + \dots \right] \right). \quad (29)$$

In light of (1) - (3), from (28) and (29), it is evident that

$$\begin{aligned} 1 + \frac{(\lambda+1)}{\gamma} \Theta_2 a_2 z + \frac{1}{\gamma} \left[(\lambda+2)\Theta_3 a_3 + \frac{(\lambda-1)(\lambda+2)}{2} \Theta_2^2 a_2^2 \right] z^2 + \dots \\ = 1 + \frac{1}{2} B_1 p_1 z + \left[\frac{1}{2} B_1 \left(p_2 - \frac{p_1^2}{2} \right) + \frac{1}{4} B_2 p_1^2 \right] z^2 + \dots \end{aligned}$$

and

$$\begin{aligned} 1 - \frac{(\lambda+1)}{\gamma} \Theta_2 a_2 w + \frac{1}{\gamma} \left[-(\lambda+2) \Theta_3 a_3 + \left(\frac{(\lambda-1)(\lambda+2)}{2} \Theta_2^2 + 2(\lambda+2) \Theta_3 \right) a_2^2 \right] w^2 + \dots \\ = 1 + \frac{1}{2} B_1 q_1 w + \left[\frac{1}{2} B_1 (q_2 - \frac{q_1^2}{2}) + \frac{1}{4} B_2 q_1^2 \right] w^2 + \dots \end{aligned}$$

which yields the following relations.

$$(1 + \lambda) \Theta_2 a_2 = \frac{\gamma}{2} B_1 p_1 \quad (30)$$

$$\left[\frac{(\lambda-1)(\lambda+2)}{2} \Theta_2^2 a_2^2 + (\lambda+2) \Theta_3 a_3 \right] = \frac{\gamma}{2} B_1 (p_2 - \frac{p_1^2}{2}) + \frac{\gamma}{4} B_2 p_1^2 \quad (31)$$

$$-(\lambda+1) \Theta_2 a_2 = \frac{\gamma}{2} B_1 q_1 \quad (32)$$

and

$$\left[\left(2(\lambda+2) \Theta_3 + \frac{(\lambda-1)(\lambda+2)}{2} \Theta_2^2 \right) a_2^2 - (\lambda+2) \Theta_3 a_3 \right] = \frac{\gamma}{2} B_1 (q_2 - \frac{q_1^2}{2}) + \frac{\gamma}{4} B_2 q_1^2. \quad (33)$$

From (30) and (32), it follows that

$$p_1 = -q_1 \quad (34)$$

and

$$8(\lambda+1)^2 \Theta_2^2 a_2^2 = \gamma^2 B_1^2 (p_1^2 + q_1^2). \quad (35)$$

Adding (31) and (33), we obtain

$$\left[(\lambda-1)(\lambda+2) \Theta_2^2 + 2(\lambda+2) \Theta_3 \right] a_2^2 = \frac{\gamma B_1}{2} (p_2 + q_2) + \frac{\gamma}{4} (B_2 - B_1) (p_1^2 + q_1^2). \quad (36)$$

Using (35) in (36), we get

$$a_2^2 = \frac{\gamma^2 B_1^3 (p_2 + q_2)}{2\gamma B_1^2 [(\lambda-1)(\lambda+2) \Theta_2^2 + 2(\lambda+2) \Theta_3] - 4(B_2 - B_1)(1+\lambda)^2 \Theta_2^2}. \quad (37)$$

Applying Lemma 5 for the coefficients p_2 and q_2 , we immediately have

$$|a_2|^2 \leq \frac{2|\gamma|^2 B_1^3}{|\gamma B_1^2 [(\lambda-1)(\lambda+2) \Theta_2^2 + 2(\lambda+2) \Theta_3] - 2(B_2 - B_1)(1+\lambda)^2 \Theta_2^2|}.$$

This gives the bound on $|a_2|$ as asserted in (22).

Next, in order to find the bound on $|a_3|$, by subtracting (33) from (31), we get

$$\left[2(\lambda+2) \Theta_3 a_3 - 2(\lambda+2) \Theta_3 a_2^2 \right] = \frac{\gamma B_1}{2} \left[(p_2 - q_2) - \frac{1}{2} (p_1^2 - q_1^2) \right] + \frac{\gamma B_2}{4} (p_1^2 - q_1^2). \quad (38)$$

Using (34) and (35) in (38), we get

$$a_3 = \frac{\gamma B_1 (p_2 - q_2)}{4(\lambda+2) \Theta_3} + \frac{\gamma^2 B_1^2 (p_1^2 + q_1^2)}{8(1+\lambda)^2 \Theta_2^2}.$$

Applying Lemma 5 once again for the coefficients p_1, q_1, p_2 and q_2 , we readily get (23). This completes the proof of Theorem 6. \square

Putting $\lambda = 0$ in Theorem 6, we have the following corollary.

Corollary 7. Let the function $f(z)$ given by (1) be in the class $\mathcal{S}_{\Sigma}^{\mu,b}(\gamma, \phi)$. Then

$$|a_2| \leq \frac{|\gamma|B_1\sqrt{B_1}}{\sqrt{|\gamma B_1^2(2\Theta_3 - \Theta_2^2) - (B_2 - B_1)\Theta_2^2|}}$$

and

$$|a_3| \leq \frac{|\gamma|B_1}{2\Theta_3} + \left(\frac{|\gamma|B_1}{\Theta_2}\right)^2.$$

Putting $\lambda = 1$ in Theorem 6, we have the following corollary.

Corollary 8. Let the function $f(z)$ given by (1) be in the class $\mathcal{H}_{\Sigma}^{\mu,b}(\gamma, \phi)$. Then

$$|a_2| \leq \frac{|\gamma|B_1\sqrt{B_1}}{\sqrt{|3\gamma B_1^2\Theta_3 - 4(B_2 - B_1)\Theta_2^2|}}$$

and

$$|a_3| \leq \frac{|\gamma|B_1}{3\Theta_3} + \left(\frac{|\gamma|B_1}{2\Theta_2}\right)^2.$$

Since $\mathcal{J}_{0,b}f(z) = f(z)$, from Corollaries 7 and 8, we get the following corollaries.

Corollary 9. Let the function $f(z)$ given by (1) be in the class $\mathcal{S}_{\Sigma}^*(\gamma, \phi)$. Then

$$|a_2| \leq \frac{|\gamma|B_1\sqrt{B_1}}{\sqrt{|\gamma B_1^2 - (B_2 - B_1)|}}$$

and

$$|a_3| \leq \frac{|\gamma|B_1}{2} + (|\gamma|B_1)^2.$$

Corollary 10. Let the function $f(z)$ given by (1) be in the class $\mathcal{H}_{\Sigma}(\gamma, \phi)$. Then

$$|a_2| \leq \frac{|\gamma|B_1\sqrt{B_1}}{\sqrt{|3\gamma B_1^2 - 4(B_2 - B_1)|}}$$

and

$$|a_3| \leq \frac{|\gamma|B_1}{3} + \left(\frac{|\gamma|B_1}{2}\right)^2.$$

Remark 11. Since $\mathcal{J}_{0,b}f(z) = f(z)$ and $\gamma = 1$, Theorem 2.1 reduces to Theorem 2.8 of Deniz [6].

3. Corollaries and Consequences

By setting $\phi(z) = \frac{1 + Az}{1 + Bz}$, $-1 \leq B < A \leq 1$, in Theorem 6, we state the following Theorem .

Theorem 12. Let the function $f(z)$ given by (1) be in the class $\mathcal{S}_{\Sigma}^{\mu,b}(\gamma, \lambda, A, B)$. Then

$$|a_2| \leq \frac{|\gamma|\sqrt{2}(A - B)}{\sqrt{|\gamma(A - B)[(\lambda - 1)(\lambda + 2)\Theta_2^2 + 2(\lambda + 2)\Theta_3] + 2(B + 1)(1 + \lambda)^2\Theta_2^2|}}$$

and

$$|a_3| \leq \frac{|\gamma|(A - B)}{(\lambda + 2)\Theta_3} + \left(\frac{|\gamma|(A - B)}{(1 + \lambda)\Theta_2}\right)^2.$$

Putting $\lambda = 0$ in Theorem 12, we have the following corollary.

Corollary 13. Let the function $f(z)$ given by (1) be in the class $\mathcal{S}_{\Sigma}^{\mu,b}(\gamma, A, B)$. Then

$$|a_2| \leq \frac{|\gamma|(A-B)}{\sqrt{|\gamma(A-B)(2\Theta_3 - \Theta_2^2) + (B+1)\Theta_2^2|}}$$

and

$$|a_3| \leq \frac{|\gamma|(A-B)}{2\Theta_3} + \left(\frac{|\gamma|(A-B)}{\Theta_2} \right)^2.$$

Putting $\lambda = 1$ in Theorem 12, we have the following corollary.

Corollary 14. Let the function $f(z)$ given by (1) be in the class $\mathcal{H}_{\Sigma}^{\mu,b}(\gamma, A, B)$. Then

$$|a_2| \leq \frac{|\gamma|(A-B)}{\sqrt{|3\gamma(A-B)\Theta_3 + 4(B+1)\Theta_2^2|}}$$

and

$$|a_3| \leq \frac{|\gamma|(A-B)}{3\Theta_3} + \left(\frac{|\gamma|(A-B)}{2\Theta_2} \right)^2.$$

Since $\mathcal{J}_{0,b}f(z) = f(z)$, from Corollaries 13 and 14, we get the following corollaries.

Corollary 15. Let the function $f(z)$ given by (1) be in the class $\mathcal{S}_{\Sigma}^*(\gamma, A, B)$. Then

$$|a_2| \leq \frac{|\gamma|(A-B)}{\sqrt{|\gamma(A-B) + (B+1)|}}$$

and

$$|a_3| \leq \frac{|\gamma|(A-B)}{2} + (|\gamma|(A-B))^2.$$

Corollary 16. Let the function $f(z)$ given by (1) be in the class $\mathcal{H}_{\Sigma}(\gamma, A, B)$. Then

$$|a_2| \leq \frac{|\gamma|(A-B)}{\sqrt{|3\gamma(A-B) + 4(B+1)|}}$$

and

$$|a_3| \leq \frac{|\gamma|(A-B)}{3} + \left(\frac{|\gamma|(A-B)}{2} \right)^2.$$

Further, by setting $\phi(z) = \frac{1+(1-2\alpha)z}{1-z}$, $0 \leq \alpha < 1$ in Theorem 6 we get the following result.

Theorem 17. Let the function $f(z)$ given by (1) be in the class $\mathcal{S}_{\Sigma}^{\mu,b}(\gamma, \lambda, \alpha)$. Then

$$|a_2| \leq \frac{2\sqrt{|\gamma|(1-\alpha)}}{\sqrt{(\lambda-1)(\lambda+2)\Theta_2^2 + 2(\lambda+2)\Theta_3}}$$

and

$$|a_3| \leq \frac{2|\gamma|(1-\alpha)}{(\lambda+2)\Theta_3} + \left(\frac{2|\gamma|(1-\alpha)}{(1+\lambda)\Theta_2} \right)^2.$$

Putting $\lambda = 0$ in Theorem 17, we have the following corollary.

Corollary 18. Let the function $f(z)$ given by (1) be in the class $\mathcal{S}_{\Sigma}^{\mu,b}(\gamma, \alpha)$. Then

$$|a_2| \leq \sqrt{\frac{2|\gamma|(1-\alpha)}{(2\Theta_3 - \Theta_2^2)}}$$

and

$$|a_3| \leq \frac{|\gamma|(1-\alpha)}{\Theta_3} + \left(\frac{2|\gamma|(1-\alpha)}{\Theta_2} \right)^2.$$

Putting $\lambda = 1$ in Theorem 17, we have the following corollary.

Corollary 19. *Let the function $f(z)$ given by (1) be in the class $\mathcal{H}_{\Sigma}^{\mu,b}(\gamma, \alpha)$. Then*

$$|a_2| \leq \sqrt{\frac{|\gamma|2(1-\alpha)}{3\Theta_3}}$$

and

$$|a_3| \leq \frac{2|\gamma|(1-\alpha)}{3\Theta_3} + \left(\frac{|\gamma|(1-\alpha)}{\Theta_2} \right)^2.$$

Since $\mathcal{J}_{0,b}f(z) = f(z)$, from Corollaries 18 and 19, we get the following corollaries.

Corollary 20. *Let the function $f(z)$ given by (1) be in the class $\mathcal{S}_{\Sigma}^*(\gamma, \alpha)$. Then*

$$|a_2| \leq \sqrt{2|\gamma|(1-\alpha)}$$

and

$$|a_3| \leq |\gamma|(1-\alpha) + 4|\gamma|^2(1-\alpha)^2.$$

Corollary 21. *Let the function $f(z)$ given by (1) be in the class $\mathcal{H}_{\Sigma}(\gamma, \alpha)$. Then*

$$|a_2| \leq \sqrt{\frac{|\gamma|2(1-\alpha)}{3}}$$

and

$$|a_3| \leq \frac{2|\gamma|(1-\alpha)}{3} + |\gamma|^2(1-\alpha)^2.$$

Concluding Remarks: By specializing the parameters μ and b , various other interesting corollaries and consequences of our main results (which are asserted by Theorem 6 above) can be derived similarly, hence we omit the details.

Acknowledgements: The authors thank the referees for their insightful suggestions.

REFERENCES

- [1] R.M. Ali, S.K. Lee, V. Ravichandran and S. Supramanian, Coefficient estimates for bi-univalent Ma-Minda starlike and convex functions, *Appl. Math. Lett.* **25** (2012), 344–351.
- [2] D. A. Brannan, J. Clunie and W. E. Kirwan, Coefficient estimates for a class of star-like functions, *Canad. J. Math.* **22** (1970), 476–485.
- [3] D.A. Brannan and J.G. Clunie (Editors), *Aspects of Contemporary Complex Analysis*, Academic Press, London, 1980.
- [4] D. A. Brannan and T. S. Taha, On some classes of bi-univalent functions, *Studia Univ. Babeş-Bolyai Math.* **31** (2) (1986), 70–77.
- [5] J. Choi and H.M. Srivastava, Certain families of sries associated with the Hurwitz-Lerch Zeta function, *Appl. Math. Comput.*, **170** (2005), 399–409.
- [6] E. Deniz, Certain subclasses of bi-univalent functions satisfying subordinate conditions, *Journal of Classical Analysis* **2**(1) (2013), 49–60.
- [7] C. Ferreira and J.L. Lopez, Asymptotic expansions of the Hurwitz-Lerch Zeta function, *J. Math. Anal. Appl.*, **298** (2004), 210–224.
- [8] T.M. Flett, The dual of an inequality of Hardy and Littlewood and some related inequalities, *J. Math. Anal. Appl.*, **38** (1972), 746–765.
- [9] B. A. Frasin and M. K. Aouf, New subclasses of bi-univalent functions, *Appl. Math. Lett.* **24** (2011), 1569–1573.

- [10] M. Garg, K. Jain and H. M. Srivastava, Some relationships between the generalized Apostol-Bernoulli polynomials and Hurwitz-Lerch Zeta functions, *Integral Transforms Spec. Funct.*, **17** (2006), 803–815.
- [11] T. Hayami and S. Owa, Coefficient bounds for bi-univalent functions, *Pan Amer. Math. J.* **22** (4) (2012), 15–26.
- [12] K. Inayat Noor, On Bazilevič functions of complex order, *Nihonkai Math. J.*, **3** (1992), 115–124.
- [13] I. B. Jung, Y. C. Kim and H. M. Srivastava, The Hardy space of analytic functions associated with certain one-parameter families of integral operators, *J. Math. Anal. Appl.*, **176** (1993), 138–147.
- [14] M. Lewin, On a coefficient problem for bi-univalent functions, *Proc. Amer. Math. Soc.* **18** (1967), 63–68.
- [15] X.-F. Li and A.-P. Wang, Two new subclasses of bi-univalent functions, *Internat. Math. Forum* **7** (2012), 1495–1504.
- [16] S.-D. Lin and H. M. Srivastava, Some families of the Hurwitz-Lerch Zeta functions and associated fractional derivative and other integral representations, *Appl. Math. comput.*, **154** (2004), 725–733.
- [17] S.-D. Lin, H.M.Srivastava and P.-Y. Wang, Some expansion formulas for a class of generalized Hurwitz-Lerch Zeta functions, *Integral Transforms Spec. Funct.*, **17** (2006), 817–827.
- [18] W.C. Ma, D. Minda, A unified treatment of some special classes of functions, in: *Proceedings of the Conference on Complex Analysis, Tianjin, 1992*, 157 - 169, Conf. Proc. Lecture Notes Anal. 1. Int. Press, Cambridge, MA, 1994.
- [19] E. Netanyahu, The minimal distance of the image boundary from the origin and the second coefficient of a univalent function in $z < 1$, *Arch. Rational Mech. Anal.* **32** (1969), 100–112.
- [20] C. Pommerenke, *Univalent Functions*, Vandenhoeck & Ruprecht, Göttingen, 1975.
- [21] J.K. Prajapat and S. P. Goyal, Applications of Srivastava-Attiya operator to the classes of strongly starlike and strongly convex functions, *J. Math. Inequal.*, **3** (2009), 129–137.
- [22] D. Răducanu and H. M. Srivastava, A new class of analytic functions defined by means of a convolution operator involving the Hurwitz-Lerch Zeta function, *Integral Transforms Spec. Funct.*, **18** (2007), 933–943
- [23] G.L. Reddy and K.S. Padmanaban, On analytic functions with reference to the Bernardi integral operator, *Bull. Austral. Math. Soc.*, **25** (1982), 387–396.
- [24] H.M. Srivastava and A. Attiya, An integral operator associated with the Hurwitz-Lerch Zeta function and differential subordination, *Integral Transforms Spec. Funct.*, **18** (2007), 207–216.
- [25] H.M. Srivastava and J. Choi, *Series associated with the Zeta and related functions*, Dordrecht, Boston, London, Kluwer Academic Publishers, 2001.
- [26] H. M. Srivastava, A. K. Mishra and P. Gochhayat, Certain subclasses of analytic and bi-univalent functions, *Appl. Math. Lett.* **23** (2010), 1188–1192.
- [27] T. S. Taha, *Topics in Univalent Function Theory*, Ph.D. Thesis, University of London, 1981.
- [28] Q.-H. Xu, Y.-C. Gui and H. M. Srivastava, Coefficient estimates for a certain subclass of analytic and bi-univalent functions, *Appl. Math. Lett.* **25** (2012), 990–994.
- [29] Q.-H. Xu, H.-G. Xiao and H. M. Srivastava, A certain general subclass of analytic and bi-univalent functions and associated coefficient estimate problems, *Appl. Math. Comput.* **218** (2012), 11461–11465.

G. MURUGUSUNDARAMOORTHY

CORRESPONDING AUTHOR, SCHOOL OF ADVANCED SCIENCES, VIT UNIVERSITY, VELLORE 632014, INDIA,

E-mail address: gmsmoorthy@yahoo.com

C. SELVARAJ

DEPARTMENT OF MATHEMATICS, PRESIDENCY COLLEGE (AUTONOMOUS), CHENNAI - 600005, INDIA

E-mail address: pamc9439@yahoo.co.in

O.S.BABU

DEPARTMENT OF MATHEMATICS, DR. AMBEDKAR GOVT. ARTS COLLEGE, CHENNAI - 600039, INDIA,

E-mail address: osbabu1009@gmail.com