

**EXISTENCE AND UNIQUENESS OF SOLUTIONS FOR
 PROBLEMS WITH WEIGHTED P-LAPLACIAN AND
 P-BIHARMONIC OPERATORS**

A. C. CAVALHEIRO

ABSTRACT. In this work we are interested in the existence and uniqueness of solutions for the Navier problem associated to the degenerate nonlinear elliptic equations

$$\begin{aligned} & \Delta(\omega(x)|\Delta u|^{p-2}\Delta u) - \sum_{j=1}^n D_j[\omega(x)\mathcal{A}_j(x, u, \nabla u)] \\ & = f_0(x) - \sum_{j=1}^n D_j f_j(x), \quad \text{in } \Omega \end{aligned}$$

in the setting of the Weighted Sobolev Spaces

1. INTRODUCTION

In this work we prove the existence and uniqueness of (weak) solutions in the weighted Sobolev space $X = W^{2,p}(\Omega, \omega) \cap W_0^{1,p}(\Omega, \omega)$ (see Definition 3) for the Navier problem

$$(P) \begin{cases} Lu(x) & = f_0(x) - \sum_{j=1}^n D_j f_j(x), \quad \text{in } \Omega \\ u(x) & = 0, \quad \text{on } \partial\Omega \\ \Delta u & = 0, \quad \text{on } \partial\Omega \end{cases}$$

where L is the partial differential operator

$$Lu(x) = \Delta(\omega(x)|\Delta u|^{p-2}\Delta u) - \sum_{j=1}^n D_j[\omega(x)\mathcal{A}_j(x, u(x), \nabla u(x))]$$

where $D_j = \partial/\partial x_j$, Ω is a bounded open set in \mathbb{R}^n , ω is a weight function, Δ is the Laplacian operator, $1 < p < \infty$ and the functions $\mathcal{A}_j : \Omega \times \mathbb{R} \times \mathbb{R}^n \rightarrow \mathbb{R}$ ($j = 1, \dots, n$) satisfies the following conditions:

- (H1)** $x \mapsto \mathcal{A}_j(x, \eta, \xi)$ is measurable on Ω for all $(\eta, \xi) \in \mathbb{R} \times \mathbb{R}^n$
 $(\eta, \xi) \mapsto \mathcal{A}_j(x, \eta, \xi)$ is continuous on $\mathbb{R} \times \mathbb{R}^n$ for almost all $x \in \Omega$.

2010 *Mathematics Subject Classification.* 35J70, 35J60.

Key words and phrases. Degenerate nonlinear equations, Weighted Sobolev spaces.

Submitted Sep. 27, 2014.

(H2) there exist a constant $\theta_1 > 0$ such that

$$[\mathcal{A}(x, \eta, \xi) - \mathcal{A}(x, \eta', \xi')].(\xi - \xi') \geq \theta_1 |\xi - \xi'|^p,$$

whenever $\xi, \xi' \in \mathbb{R}^n$, $\xi \neq \xi'$, where $\mathcal{A}(x, \eta, \xi) = (\mathcal{A}_1(x, \eta, \xi), \dots, \mathcal{A}_n(x, \eta, \xi))$ (where a dot denote here the Euclidian scalar product in \mathbb{R}^n).

(H3) $\mathcal{A}(x, \eta, \xi). \xi \geq \lambda_1 |\xi|^p$, where λ_1 is a positive constant.

(H4) $|\mathcal{A}(x, \eta, \xi)| \leq K_1(x) + h_1(x)|\eta|^{p/p'} + h_2(x)|\xi|^{p/p'}$, where K_1, h_1 and h_2 are positive functions, with h_1 and $h_2 \in L^\infty(\Omega)$, and $K_1 \in L^{p'}(\Omega, \omega)$ (with $1/p + 1/p' = 1$).

By a *weight*, we shall mean a locally integrable function ω on \mathbb{R}^n such that $\omega(x) > 0$ for a.e. $x \in \mathbb{R}^n$. Every weight ω gives rise to a measure on the measurable subsets on \mathbb{R}^n through integration. This measure will be denoted by μ . Thus, $\mu(E) = \int_E \omega(x) dx$ for measurable sets $E \subset \mathbb{R}^n$.

In general, the Sobolev spaces $W^{k,p}(\Omega)$ without weights occur as spaces of solutions for elliptic and parabolic partial differential equations. For degenerate partial differential equations, i.e., equations with various types of singularities in the coefficients, it is natural to look for solutions in weighted Sobolev spaces (see [1], [2] and [4]).

In various applications, we can meet boundary value problems for elliptic equations whose ellipticity is disturbed in the sense that some degeneration or singularity appears. This bad behaviour can be caused by the coefficients of the corresponding differential operator as well as by the solution itself. The so-called p-Laplacian is a prototype of such an operator and its character can be interpreted as a degeneration or as a singularity of the classical (linear) Laplace operator (with $p = 2$). There are several very concrete problems from practice which lead to such differential equations, e.g. from glaciology, non-Newtonian fluid mechanics, flows through porous media, differential geometry, celestial mechanics, climatology, petroleum extraction, reaction-diffusion problems, etc.

A class of weights, which is particularly well understood, is the class of A_p -weights (or Muckenhoupt class) that was introduced by B. Muckenhoupt (see [11]). These classes have found many useful applications in harmonic analysis (see [13]). Another reason for studying A_p -weights is the fact that powers of the distance to submanifolds of \mathbb{R}^n often belong to A_p (see [10]). There are, in fact, many interesting examples of weights (see [9] for p-admissible weights).

In the non-degenerate case (i.e. with $\omega(x) \equiv 1$), for all $f \in L^p(\Omega)$ the Poisson equation associated with the Dirichlet problem

$$\begin{cases} -\Delta u = f(x), & \text{in } \Omega \\ u(x) = 0, & \text{on } \partial\Omega \end{cases}$$

is uniquely solvable in $W^{2,p}(\Omega) \cap W_0^{1,p}(\Omega)$ (see [8]), and the nonlinear Dirichlet problem

$$\begin{cases} -\Delta_p u = f(x), & \text{in } \Omega \\ u(x) = 0, & \text{on } \partial\Omega \end{cases}$$

is uniquely solvable in $W_0^{1,p}(\Omega)$ (see [3]), where $\Delta_p u = \operatorname{div}(|\nabla u|^{p-2} \nabla u)$ is the p-Laplacian operator. In the degenerate case, the weighted p-Biharmonic operator

have been studied by many authors (see [12] and the references therein), and the degenerated p-Laplacian has been studied in [4].

The following theorem will be proved in section 3.

Theorem 1 Assume (H1)-(H4). If $\omega \in A_p$ (with $1 < p < \infty$), $f_j/\omega \in L^{p'}(\Omega, \omega)$ ($j = 0, 1, \dots, n$ and $1/p + 1/p' = 1$) then the problem (P) has a unique solution $u \in X = W^{2,p}(\Omega, \omega) \cap W_0^{1,p}(\Omega, \omega)$. Moreover, we have

$$\|u\|_X \leq \frac{1}{\gamma^{p'/p}} \left(C_\Omega \|f_0/\omega\|_{L^{p'}(\Omega, \omega)} + \sum_{j=1}^n \|f_j/\omega\|_{L^{p'}(\Omega, \omega)} \right)^{p'/p},$$

where $\gamma = \min \{\lambda_1, 1\}$.

2. DEFINITIONS AND BASIC RESULTS

Let ω be a locally integrable nonnegative function in \mathbb{R}^n and assume that $0 < \omega(x) < \infty$ almost everywhere. We say that ω belongs to the Muckenhoupt class A_p , $1 < p < \infty$, or that ω is an A_p -weight, if there is a constant $C = C_{p,\omega}$ such that

$$\left(\frac{1}{|B|} \int_B \omega(x) dx \right) \left(\frac{1}{|B|} \int_B \omega^{1/(1-p)}(x) dx \right)^{p-1} \leq C$$

for all balls $B \subset \mathbb{R}^n$, where $|\cdot|$ denotes the n -dimensional Lebesgue measure in \mathbb{R}^n . If $1 < q \leq p$, then $A_q \subset A_p$ (see [7],[9] or [13] for more information about A_p -weights). The weight ω satisfies the doubling condition if there exists a positive constant C such that $\mu(B(x; 2r)) \leq C \mu(B(x; r))$ for every ball $B = B(x; r) \subset \mathbb{R}^n$, where $\mu(B) = \int_B \omega(x) dx$. If $\omega \in A_p$, then μ is doubling (see Corollary 15.7 in [9]).

As an example of A_p -weight, the function $\omega(x) = |x|^\alpha$, $x \in \mathbb{R}^n$, is in A_p if and only if $-n < \alpha < n(p-1)$ (see Corollary 4.4, Chapter IX in [13]).

If $\omega \in A_p$, then $\left(\frac{|E|}{|B|} \right)^p \leq C \frac{\mu(E)}{\mu(B)}$ whenever B is a ball in \mathbb{R}^n and E is a measurable subset of B (see 15.5 *strong doubling property* in [9]). Therefore, if $\mu(E) = 0$ then $|E| = 0$.

Definition 1 Let ω be a weight, and let $\Omega \subset \mathbb{R}^n$ be open. For $1 < p < \infty$ we define $L^p(\Omega, \omega)$ as the set of measurable functions f on Ω such that

$$\|f\|_{L^p(\Omega, \omega)} = \left(\int_\Omega |f(x)|^p \omega(x) dx \right)^{1/p} < \infty.$$

If $\omega \in A_p$, $1 < p < \infty$, then $\omega^{-1/(p-1)}$ is locally integrable and we have $L^p(\Omega, \omega) \subset L_{\text{loc}}^1(\Omega)$ for every open set Ω (see Remark 1.2.4 in [14]). It thus makes sense to talk about weak derivatives of functions in $L^p(\Omega, \omega)$.

Definition 2 Let $\Omega \subset \mathbb{R}^n$ be open and let $\omega \in A_p$ ($1 < p < \infty$). We define the weighted Sobolev space $W^{1,p}(\Omega, \omega)$ as the set of functions $u \in L^p(\Omega, \omega)$ with weak derivatives $D_j u \in L^p(\Omega, \omega)$ for $j = 1, \dots, n$. The norm of u in $W^{1,p}(\Omega, \omega)$ is defined by

$$\|u\|_{W^{1,p}(\Omega, \omega)} = \left(\int_\Omega |u(x)|^p \omega(x) dx + \sum_{j=1}^n \int_\Omega |D_j u(x)|^p \omega(x) dx \right)^{1/p}. \quad (1)$$

We also define $W_0^{1,p}(\Omega, \omega)$ as the closure of $C_0^\infty(\Omega)$ with respect to the norm $\|\cdot\|_{W^{1,p}(\Omega, \omega)}$.

If $\omega \in A_p$, then $W^{1,p}(\Omega, \omega)$ is the closure of $C^\infty(\Omega)$ with respect to the norm (1) (see Theorem 2.1.4 in [14]). The spaces $W^{1,p}(\Omega, \omega)$ and $W_0^{1,p}(\Omega, \omega)$ are Banach spaces.

It is evident that the weight function ω which satisfy $0 < c_1 \leq \omega(x) \leq c_2$ for $x \in \Omega$ (c_1 and c_2 positive constants), give nothing new (the space $W_0^{1,p}(\Omega, \omega)$ is then identical with the classical Sobolev space $W_0^{1,p}(\Omega)$). Consequently, we shall interested above all in such weight functions ω which either vanish somewhere in Ω or increase to infinity (or both).

In this article we use the following results.

Theorem 2 Let $\omega \in A_p$, $1 < p < \infty$, and let Ω be a bounded open set in \mathbb{R}^n . If $u_m \rightarrow u$ in $L^p(\Omega, \omega)$ then there exist a subsequence $\{u_{m_k}\}$ and a function $\Phi \in L^p(\Omega, \omega)$ such that

- (i) $u_{m_k}(x) \rightarrow u(x)$, $m_k \rightarrow \infty$, μ -a.e. on Ω ;
 - (ii) $|u_{m_k}(x)| \leq \Phi(x)$, μ -a.e. on Ω ;
- (where $\mu(E) = \int_E \omega(x) dx$).

Proof. The proof of this theorem follows the lines of Theorem 2.8.1 in [6]. \square

Theorem 3 (The weighted Sobolev inequality) Let Ω be an open bounded set in \mathbb{R}^n and $\omega \in A_p$ ($1 < p < \infty$). There exist positive constants C_Ω and δ such that for all $u \in W_0^{1,p}(\Omega, \omega)$ and all k satisfying $1 \leq k \leq n/(n-1) + \delta$,

$$\|u\|_{L^{kp}(\Omega, \omega)} \leq C_\Omega \|\nabla u\|_{L^p(\Omega, \omega)}. \quad (2)$$

Proof. Its suffices to prove the inequality for functions $u \in C_0^\infty(\Omega)$ (see Theorem 1.3 in [5]). To extend the estimates (2) to arbitrary $u \in W_0^{1,p}(\Omega, \omega)$, we let $\{u_m\}$ be a sequence of $C_0^\infty(\Omega)$ functions tending to u in $W_0^{1,p}(\Omega, \omega)$. Applying the estimates (2) to differences $u_{m_1} - u_{m_2}$, we see that $\{u_m\}$ will be a Cauchy sequence in $L^{kp}(\Omega, \omega)$. Consequently the limit function u will lie in the desired spaces and satisfy (2). \square

Lemma 1 Let $1 < p < \infty$.

(a) There exists a constant α_p such that

$$\left| |x|^{p-2}x - |y|^{p-2}y \right| \leq \alpha_p |x - y| (|x| + |y|)^{p-2},$$

for all $x, y \in \mathbb{R}^n$;

(b) There exist two positive constants β_p, γ_p such that for every $x, y \in \mathbb{R}^n$

$$\beta_p (|x| + |y|)^{p-2} |x - y|^2 \leq (|x|^{p-2}x - |y|^{p-2}y) \cdot (x - y) \leq \gamma_p (|x| + |y|)^{p-2} |x - y|^2.$$

Proof. See [3], Proposition 17.2 and Proposition 17.3. \square

Definition 3 We say that an element $u \in X = W^{2,p}(\Omega, \omega) \cap W_0^{1,p}(\Omega, \omega)$ is a (weak) solution of problem (P) if, for all $\varphi \in X$,

$$\begin{aligned} & \int_{\Omega} |\Delta u|^{p-2} \Delta u \Delta \varphi \omega dx + \sum_{j=1}^n \int_{\Omega} \omega(x) \mathcal{A}_j(x, u(x), \nabla u(x)) D_j \varphi(x) dx \\ & = \int_{\Omega} f_0(x) \varphi(x) dx + \sum_{j=1}^n \int_{\Omega} f_j(x) D_j \varphi(x) dx, \end{aligned}$$

3. PROOF OF THEOREM 1

The basic idea is to reduce the problem (P) to an operator equation $Au = T$ and apply the theorem below.

Theorem 4 Let $A : X \rightarrow X^*$ be a monotone, coercive and hemicontinuous operator on the real, separable, reflexive Banach space X . Then the following assertions hold:

- (a) For each $T \in X^*$ the equation $Au = T$ has a solution $u \in X$;
- (b) If the operator A is strictly monotone, then equation $Au = T$ is uniquely solvable in X .

Proof. See Theorem 26.A in [16]. □

To prove the existence of solutions, we define $B, B_1, B_2 : X \times X \rightarrow \mathbb{R}$ and $T : X \rightarrow \mathbb{R}$ by

$$\begin{aligned}
 B(u, \varphi) &= B_1(u, \varphi) + B_2(u, \varphi) \\
 B_1(u, \varphi) &= \sum_{j=1}^n \int_{\Omega} \omega \mathcal{A}_j(x, u, \nabla u) D_j \varphi dx = \int_{\Omega} \omega \mathcal{A}(x, u, \nabla u) \cdot \nabla \varphi dx \\
 B_2(u, \varphi) &= \int_{\Omega} |\Delta u|^{p-2} \Delta u \Delta \varphi \omega dx \\
 T(\varphi) &= \int_{\Omega} f_0(x) \varphi(x) dx + \sum_{j=1}^n \int_{\Omega} f_j(x) D_j \varphi(x) dx.
 \end{aligned}$$

Then $u \in X$ is a (weak) solution to problem (P) if

$$B(u, \varphi) = B_1(u, \varphi) + B_2(u, \varphi) = T(\varphi),$$

for all $\varphi \in X$.

Step 1 For $j = 1, \dots, n$ we define the operator $F_j : X \rightarrow L^{p'}(\Omega, \omega)$ by

$$(F_j u)(x) = \mathcal{A}_j(x, u(x), \nabla u(x)).$$

We have that the operator F_j is bounded and continuous. In fact:

(i) Using (H4) we obtain

$$\begin{aligned}
 \|F_j u\|_{L^{p'}(\Omega, \omega)}^{p'} &= \int_{\Omega} |F_j u(x)|^{p'} \omega dx = \int_{\Omega} |\mathcal{A}_j(x, u, \nabla u)|^{p'} \omega dx \\
 &\leq \int_{\Omega} \left(K_1 + h_1 |u|^{p/p'} + h_2 |\nabla u|^{p/p'} \right)^{p'} \omega dx \\
 &\leq C_p \int_{\Omega} \left[(K_1^{p'} + h_1^{p'} |u|^p + h_2^{p'} |\nabla u|^p) \omega \right] dx \\
 &= C_p \left[\int_{\Omega} K_1^{p'} \omega dx + \int_{\Omega} h_1^{p'} |u|^p \omega dx + \int_{\Omega} h_2^{p'} |\nabla u|^p \omega dx \right], \tag{3}
 \end{aligned}$$

where the constant C_p depends only on p . We have, by Theorem 3,

$$\begin{aligned}
 \int_{\Omega} h_1^{p'} |u|^p \omega dx &\leq \|h_1\|_{L^\infty(\Omega)}^{p'} \int_{\Omega} |u|^p \omega dx \\
 &\leq C_\Omega^p \|h_1\|_{L^\infty(\Omega)}^{p'} \int_{\Omega} |\nabla u|^p \omega dx \\
 &\leq C_\Omega^p \|h_1\|_{L^\infty(\Omega)}^{p'} \|u\|_X^p,
 \end{aligned}$$

and $\int_{\Omega} h_2^{p'} |\nabla u|^p \omega dx \leq \|h_2\|_{L^\infty(\Omega)}^{p'} \int_{\Omega} |\nabla u|^p \omega dx \leq \|h_2\|_{L^\infty(\Omega)}^{p'} \|u\|_X^p$. Therefore, in (3) we obtain

$$\|F_j u\|_{L^{p'}(\Omega, \omega)} \leq C_p \left(\|K\|_{L^{p'}(\Omega, \omega)} + (C_\Omega^{p/p'}) \|h_1\|_{L^\infty(\Omega)} + \|h_2\|_{L^\infty(\Omega)} \|u\|_X^{p/p'} \right).$$

(ii) Let $u_m \rightarrow u$ in X as $m \rightarrow \infty$. We need to show that $F_j u_m \rightarrow F_j u$ in $L^{p'}(\Omega, \omega)$.

If $u_m \rightarrow u$ in X , then $u_m \rightarrow u$ in $L^p(\Omega, \omega)$ and $|\nabla u_m| \rightarrow |\nabla u|$ in $L^p(\Omega, \omega)$. Using Theorem 2, there exist a subsequence $\{u_{m_k}\}$ and functions Φ_1 and Φ_2 in $L^p(\Omega, \omega)$ such that

$$\begin{aligned} u_{m_k}(x) &\rightarrow u(x), \quad \mu - \text{a.e. in } \Omega, \\ |u_{m_k}(x)| &\leq \Phi_1(x), \quad \mu - \text{a.e. in } \Omega, \\ |\nabla u_{m_k}(x)| &\rightarrow |\nabla u(x)|, \quad \mu - \text{a.e. in } \Omega, \\ |\nabla u_{m_k}(x)| &\leq \Phi_2(x), \quad \mu - \text{a.e. in } \Omega. \end{aligned}$$

Hence, using (H4), we obtain

$$\begin{aligned} \|F_j u_{m_k} - F_j u\|_{L^{p'}(\Omega, \omega)}^{p'} &= \int_{\Omega} |F_j u_{m_k}(x) - F_j u(x)|^{p'} \omega dx \\ &= \int_{\Omega} |\mathcal{A}_j(x, u_{m_k}, \nabla u_{m_k}) - \mathcal{A}_j(x, u, \nabla u)|^{p'} \omega dx \\ &\leq C_p \int_{\Omega} \left(|\mathcal{A}_j(x, u_{m_k}, \nabla u_{m_k})|^{p'} + |\mathcal{A}_j(x, u, \nabla u)|^{p'} \right) \omega dx \\ &\leq C_p \left[\int_{\Omega} \left(K_1 + h_1 |u_{m_k}|^{p/p'} + h_2 |\nabla u_{m_k}|^{p/p'} \right)^{p'} \omega dx \right. \\ &\quad \left. + \int_{\Omega} \left(K_1 + h_1 |u|^{p/p'} + h_2 |\nabla u|^{p/p'} \right)^{p'} \omega dx \right] \\ &\leq 2 C_p \int_{\Omega} \left(K_1 + h_1 \Phi_1^{p/p'} + h_2 \Phi_2^{p/p'} \right)^{p'} \omega dx \\ &\leq 2 C_p \left[\int_{\Omega} K_1^{p'} \omega dx + \int_{\Omega} h_1^{p'} \Phi_1^p \omega dx + \int_{\Omega} h_2^{p'} \Phi_2^p \omega dx \right] \\ &\leq 2 C_p \left[\|K_1\|_{L^{p'}(\Omega, \omega)}^{p'} + \|h_1\|_{L^\infty(\Omega)}^{p'} \int_{\Omega} \Phi_1^p \omega dx \right. \\ &\quad \left. + \|h_2\|_{L^\infty(\Omega)}^{p'} \int_{\Omega} \Phi_2^p \omega dx \right] \\ &\leq 2 C_p \left[\|K_1\|_{L^{p'}(\Omega, \omega)}^{p'} + \|h_1\|_{L^\infty(\Omega)}^{p'} \|\Phi_1\|_{L^p(\Omega, \omega)}^p \right. \\ &\quad \left. + \|h_2\|_{L^\infty(\Omega)}^{p'} \|\Phi_2\|_{L^p(\Omega, \omega)}^p \right]. \end{aligned}$$

By condition (H1), we have

$$F_j u_m(x) = \mathcal{A}_j(x, u_m(x), \nabla u_m(x)) \rightarrow \mathcal{A}_j(x, u(x), \nabla u(x)) = F_j u(x),$$

as $m \rightarrow +\infty$. Therefore, by the Dominated Convergence Theorem, we obtain

$$\|F_j u_{m_k} - F_j u\|_{L^{p'}(\Omega, \omega)} \rightarrow 0,$$

that is,

$$F_j u_{m_k} \rightarrow F_j u \text{ in } L^{p'}(\Omega, \omega).$$

By the Convergence Principle in Banach spaces (see Proposition 10.13 in [15]), we have

$$F_j u_m \rightarrow F_j u \text{ in } L^{p'}(\Omega, \omega). \quad (4)$$

Step 2 We define the operator

$$\begin{aligned} G : X &\rightarrow L^{p'}(\Omega, \omega) \\ (Gu)(x) &= |\Delta u(x)|^{p-2} \Delta u(x). \end{aligned}$$

We also have that the operator G is continuous and bounded. In fact:

(i) We have

$$\begin{aligned} \|Gu\|_{L^{p'}(\Omega, \omega)}^{p'} &= \int_{\Omega} |\Delta u|^{p-2} \Delta u|^{p'} \omega dx \\ &= \int_{\Omega} |\Delta u|^{(p-2)p'} |\Delta u|^{p'} \omega dx \\ &= \int_{\Omega} |\Delta u|^p \omega dx \\ &\leq \|u\|_X^p. \end{aligned}$$

Hence, $\|Gu\|_{L^{p'}(\Omega, \omega)} \leq \|u\|_X^{p/p'}$.

(ii) If $u_m \rightarrow u$ in X then $\Delta u_m \rightarrow \Delta u$ in $L^p(\Omega, \omega)$. By Theorem 2, there exist a subsequence $\{u_{m_k}\}$ and a function $\Phi_3 \in L^p(\Omega, \omega)$ such that

$$\begin{aligned} \Delta u_{m_k}(x) &\rightarrow \Delta u(x), \quad \mu - a.e. \text{ in } \Omega \\ |\Delta u_{m_k}(x)| &\leq \Phi_3(x), \quad \mu - a.e. \text{ in } \Omega. \end{aligned}$$

Hence, using Lemma 1(a), we obtain, if $p \neq 2$

$$\begin{aligned} \|Gu_{m_k} - Gu\|_{L^{p'}(\Omega, \omega)}^{p'} &= \int_{\Omega} |Gu_{m_k} - Gu|^{p'} \omega dx \\ &= \int_{\Omega} \left| |\Delta u_{m_k}|^{p-2} \Delta u_{m_k} - |\Delta u|^{p-2} \Delta u \right|^{p'} \omega dx \\ &\leq \int_{\Omega} \left[\alpha_p |\Delta u_{m_k} - \Delta u| (|\Delta u_{m_k}| + |\Delta u|)^{(p-2)} \right]^{p'} \omega dx \\ &\leq \alpha_p^{p'} \int_{\Omega} |\Delta u_{m_k} - \Delta u|^{p'} (2\Phi_3)^{(p-2)p'} \omega dx \\ &\leq \alpha_p^{p'} 2^{(p-2)p'} \left(\int_{\Omega} |\Delta u_{m_k} - \Delta u|^p \omega dx \right)^{p'/p} \times \\ &\quad \times \left(\int_{\Omega} \Phi_3^{(p-2)p p'/(p-p')} \omega dx \right)^{(p-p')/p} \\ &\leq \alpha_p^{p'} 2^{(p-2)p'} \|u_{m_k} - u\|_X^{p'} \|\Phi\|_{L^p(\Omega, \omega)}^{p-p'}, \end{aligned}$$

since $(p-2)p p'/(p-p') = p$ if $p \neq 2$. If $p = 2$, we have

$$\begin{aligned} \|Gu_{m_k} - Gu\|_{L^2(\Omega, \omega)}^2 &=? \int_{\Omega} |\Delta u_{m_k} - \Delta u|^2 \omega dx \\ &\leq \|u_{m_k} - u\|_X^2. \end{aligned}$$

Therefore (for $1 < p < \infty$), by the Dominated Convergence Theorem, we obtain

$$\|Gu_{m_k} - Gu\|_X \rightarrow 0,$$

that is, $Gu_{m_k} \rightarrow Gu$ in $L^{p'}(\Omega, \omega)$. By convergence principle in Banach spaces (see Proposition 10.13 in [15]), we have

$$Gu_m \rightarrow Gu \text{ in } L^{p'}(\Omega, \omega). \quad (5)$$

Step 3 We have, by Theorem 3,

$$\begin{aligned} |T(\varphi)| &\leq \int_{\Omega} |f_0| |\varphi| dx + \sum_{j=1}^n \int_{\Omega} |f_j| |D_j \varphi| dx \\ &= \int_{\Omega} \frac{|f_0|}{\omega} |\varphi| \omega dx + \sum_{j=1}^n \int_{\Omega} \frac{|f_j|}{\omega} |D_j \varphi| \omega dx \\ &\leq \|f_0/\omega\|_{L^{p'}(\Omega, \omega)} \|\varphi\|_{L^p(\Omega, \omega)} + \sum_{j=1}^n \|f_j/\omega\|_{L^{p'}(\Omega, \omega)} \|D_j \varphi\|_{L^p(\Omega, \omega)} \\ &\leq \left(C_{\Omega} \|f_0/\omega\|_{L^{p'}(\Omega, \omega)} + \sum_{j=1}^n \|f_j/\omega\|_{L^{p'}(\Omega, \omega)} \right) \|\varphi\|_X. \end{aligned}$$

Moreover, using (H4) and the Hölder inequality, we also have

$$\begin{aligned} |B(u, \varphi)| &\leq |B_1(u, \varphi)| + |B_2(u, \varphi)| \\ &\leq \sum_{j=1}^n \int_{\Omega} |\mathcal{A}_j(x, u, \nabla u)| |D_j \varphi| \omega dx + \int_{\Omega} |\Delta u|^{p-2} |\Delta u| |\Delta \varphi| \omega dx. \quad (6) \end{aligned}$$

In (6) we have

$$\begin{aligned} \int_{\Omega} |\mathcal{A}(x, u, \nabla u)| |\nabla \varphi| \omega dx &\leq \int_{\Omega} \left(K_1 + h_1 |u|^{p/p'} + h_2 |\nabla u|^{p/p'} \right) |\nabla \varphi| \omega dx \\ &\leq \|K_1\|_{L^{p'}(\Omega, \omega)} \|\nabla \varphi\|_{L^p(\Omega, \omega)} + \|h_1\|_{L^{\infty}(\Omega)} \|u\|_{L^{p/p'}(\Omega, \omega)}^{p/p'} \|\nabla \varphi\|_{L^p(\Omega, \omega)} \\ &+ \|h_2\|_{L^{\infty}(\Omega)} \|\nabla u\|_{L^{p/p'}(\Omega, \omega)}^{p/p'} \|\nabla \varphi\|_{L^p(\Omega, \omega)} \\ &\leq \left(\|K_1\|_{L^{p'}(\Omega, \omega)} + (C_{\Omega}^{p/p'} \|h_1\|_{L^{\infty}(\Omega)} + \|h_2\|_{L^{\infty}(\Omega)}) \|u\|_X^{p/p'} \right) \|\varphi\|_X, \end{aligned}$$

and

$$\begin{aligned} \int_{\Omega} |\Delta u|^{p-2} |\Delta u| |\Delta \varphi| \omega dx &= \int_{\Omega} |\Delta u|^{p-1} |\Delta \varphi| \omega dx \\ &\leq \left(\int_{\Omega} |\Delta u|^p \omega dx \right)^{1/p'} \left(\int_{\Omega} |\Delta \varphi|^p \omega dx \right)^{1/p} \\ &\leq \|u\|_X^{p/p'} \|\varphi\|_X. \end{aligned}$$

Therefore, in (6) we obtain, for all $u, \varphi \in X$

$$\begin{aligned} |B(u, \varphi)| &\leq \left[\|K_1\|_{L^{p'}(\Omega, \omega)} + C_{\Omega}^{p/p'} \|h_1\|_{L^{\infty}(\Omega)} \|u\|_X^{p/p'} \right. \\ &\quad \left. + \|h_2\|_{L^{\infty}(\Omega, \omega)} \|u\|_X^{p/p'} + \|u\|_X^{p/p'} \right] \|\varphi\|_X. \end{aligned}$$

Since $B(u, \cdot)$ is linear, for each $u \in X$, there exists a linear and continuous operator $A : X \rightarrow X^*$ such that $\langle Au, \varphi \rangle = B(u, \varphi)$, for all $u, \varphi \in X$ (where $\langle f, x \rangle$ denotes the value of the linear functional f at the point x) and

$$\|Au\|_* \leq \|K_1\|_{L^{p'}(\Omega, \omega)} + C_\Omega^{p/p'} \|h_1\|_{L^\infty(\Omega)} \|u\|_X^{p/p'} + \|h_2\|_{L^\infty(\Omega, \omega)} \|u\|_X^{p/p'} + \|u\|_X^{p/p'}.$$

Consequently, problem (P) is equivalent to the operator equation

$$Au = T, \quad u \in X.$$

Step 4 Using condition (H2) and Lemma 1(b), we have

$$\begin{aligned} \langle Au_1 - Au_2, u_1 - u_2 \rangle &= B(u_1, u_1 - u_2) - B(u_2, u_1 - u_2) \\ &= \int_\Omega \omega \mathcal{A}(x, u_1, \nabla u_1) \cdot \nabla(u_1 - u_2) \, dx + \int_\Omega |\Delta u_1|^{p-2} \Delta u_1 \Delta(u_1 - u_2) \omega \, dx \\ &\quad - \int_\Omega \omega \mathcal{A}(x, u_2, \nabla u_2) \cdot \nabla(u_1 - u_2) \, dx - \int_\Omega |\Delta u_2|^{p-2} \Delta u_2 \Delta(u_1 - u_2) \omega \, dx \\ &= \int_\Omega \omega \left(\mathcal{A}(x, u_1, \nabla u_1) - \mathcal{A}(x, u_2, \nabla u_2) \right) \cdot \nabla(u_1 - u_2) \, dx \\ &\quad + \int_\Omega (|\Delta u_1|^{p-2} \Delta u_1 - |\Delta u_2|^{p-2} \Delta u_2) \Delta(u_1 - u_2) \omega \, dx \\ &\geq \theta_1 \int_\Omega \omega |\nabla(u_1 - u_2)|^p \, dx + \beta_p \int_\Omega (|\Delta u_1| + |\Delta u_2|)^{p-2} |\Delta u_1 - \Delta u_2|^2 \omega \, dx \\ &\geq \theta_1 \int_\Omega \omega |\nabla(u_1 - u_2)|^p \, dx + \beta_p \int_\Omega (|\Delta u_1 - \Delta u_2|)^{p-2} |\Delta u_1 - \Delta u_2|^2 \omega \, dx \\ &= \theta_1 \int_\Omega \omega |\nabla(u_1 - u_2)|^p \, dx + \beta_p \int_\Omega |\Delta u_1 - \Delta u_2|^p \omega \, dx \\ &\geq \theta \|u_1 - u_2\|_X^p \end{aligned}$$

where $\theta = \min \{\theta_1, \beta_p\}$.

Therefore, the operator A is strictly monotone. Moreover, using (H3), we obtain

$$\begin{aligned} \langle Au, u \rangle &= B(u, u) = B_1(u, u) + B_2(u, u) \\ &= \int_\Omega \omega \mathcal{A}(x, u, \nabla u) \cdot \nabla u \, dx + \int_\Omega |\Delta u|^{p-2} \Delta u \Delta u \omega \, dx \\ &\geq \int_\Omega \lambda_1 |\nabla u|^p \omega \, dx + \int_\Omega |\Delta u|^p \omega \, dx \\ &\geq \gamma \|u\|_X^p \end{aligned}$$

where $\gamma = \min \{\lambda_1, 1\}$. Hence, since $p > 1$, we have

$$\frac{\langle Au, u \rangle}{\|u\|_X} \rightarrow +\infty, \quad \text{as } \|u\|_X \rightarrow +\infty,$$

that is, A is coercive.

Step 5 We show that the operator A is continuous which, in particular means that A is hemicontinuous.

Let $u_m \rightarrow u$ in X as $m \rightarrow \infty$. We have,

$$\begin{aligned} |B_1(u_m, \varphi) - B_1(u, \varphi)| &\leq \sum_{j=1}^n \int_{\Omega} |\mathcal{A}_j(x, u_m, \nabla u_m) - \mathcal{A}_j(x, u, \nabla u)| |D_j \varphi| \omega \, dx \\ &= \sum_{j=1}^n \int_{\Omega} |F_j u_m - F_j u| |D_j \varphi| \omega \, dx \\ &\leq \sum_{j=1}^n \|F_j u_m - F_j u\|_{L^{p'}(\Omega, \omega)} \|D_j \varphi\|_{L^p(\Omega, \omega)} \\ &\leq \sum_{j=1}^n \|F_j u_m - F_j u\|_{L^{p'}(\Omega, \omega)} \|\varphi\|_X, \end{aligned}$$

and

$$\begin{aligned} &|B_2(u_m, \varphi) - B_2(u, \varphi)| \\ &= \left| \int_{\Omega} |\Delta u_m|^{p-2} \Delta u_m \Delta \varphi \omega \, dx - \int_{\Omega} |\Delta u|^{p-2} \Delta u \Delta \varphi \omega \, dx \right| \\ &\leq \int_{\Omega} \left| |\Delta u_m|^{p-2} \Delta u_m - |\Delta u|^{p-2} \Delta u \right| |\Delta \varphi| \omega \, dx \\ &= \int_{\Omega} |G u_m - G u| |\Delta \varphi| \omega \, dx \\ &\leq \|G u_m - G u\|_{L^{p'}(\Omega, \omega)} \|\varphi\|_X. \end{aligned}$$

for all $\varphi \in X$. Hence,

$$\begin{aligned} |B(u_m, \varphi) - B(u, \varphi)| &\leq |B_1(u_m, \varphi) - B_1(u, \varphi)| + |B_2(u_m, \varphi) - B_2(u, \varphi)| \\ &\leq \left[\sum_{j=1}^n \|F_j u_m - F_j u\|_{L^{p'}(\Omega, \omega)} + \|G u_m - G u\|_{L^{p'}(\Omega, \omega)} \right] \|\varphi\|_X. \end{aligned}$$

Then we obtain

$$\|A u_m - A u\|_* \leq \sum_{j=1}^n \|F_j u_m - F_j u\|_{L^{p'}(\Omega, \omega)} + \|G u_m - G u\|_{L^{p'}(\Omega, \omega)}.$$

Therefore, using (4) and (5) we have $\|A u_m - A u\|_* \rightarrow 0$ as $m \rightarrow +\infty$, that is, A is continuous.

Therefore, by Theorem 4, the operator equation $Au = T$ has a unique solution $u \in X$ and it is the unique solution for problem (P).

Step 6 In particular, by setting $\varphi = u$ in Definition 3, we have

$$B(u, u) = B_1(u, u) + B_2(u, u) = T(u). \quad (7)$$

Hence, using (H3) and $\gamma = \min \{\lambda_1, 1\}$, we obtain

$$\begin{aligned} B_1(u, u) + B_2(u, u) &= \int_{\Omega} \omega \mathcal{A}(x, u, \nabla u) \cdot \nabla u \, dx + \int_{\Omega} |\Delta u|^{p-2} \Delta u \Delta u \omega \, dx \\ &\geq \int_{\Omega} \lambda_1 |\nabla u|^p + \int_{\Omega} |\Delta u|^p \omega \, dx \\ &\geq \gamma \|u\|_X^p \end{aligned}$$

and

$$\begin{aligned} T(u) &= \int_{\Omega} f_0 u \, dx + \sum_{j=1}^n \int_{\Omega} f_j D_j u \, dx \\ &\leq \|f_0/\omega\|_{L^{p'}(\Omega, \omega)} \|u\|_{L^p(\Omega, \omega)} + \sum_{j=1}^n \|f_j/\omega\|_{L^{p'}(\Omega)} \|D_j u\|_{L^p(\Omega, \omega)} \\ &\leq \left(C_{\Omega} \|f_0/\omega\|_{L^{p'}(\Omega, \omega)} + \sum_{j=1}^n \|f_j/\omega\|_{L^{p'}(\Omega)} \right) \|u\|_X. \end{aligned}$$

Therefore, in (7), we have

$$\gamma \|u\|_X^p \leq \left(C_{\Omega} \|f_0/\omega\|_{L^{p'}(\Omega, \omega)} + \sum_{j=1}^n \|f_j/\omega\|_{L^{p'}(\Omega, \omega)} \right) \|u\|_X,$$

and we obtain

$$\|u\|_X \leq \frac{1}{\gamma^{p'/p}} \left(C_{\Omega} \|f_0/\omega\|_{L^{p'}(\Omega, \omega)} + \sum_{j=1}^n \|f_j/\omega\|_{L^{p'}(\Omega, \omega)} \right)^{p'/p}.$$

Example Let $\Omega = \{(x, y) \in \mathbb{R}^2 : x^2 + y^2 < 1\}$, the weight function $\omega(x, y) = (x^2 + y^2)^{-1/2}$ ($\omega \in A_3$, $p = 3$), and the function

$$\begin{aligned} \mathcal{A} : \Omega \times \times \mathbb{R}^2 &\rightarrow \mathbb{R}^2 \\ \mathcal{A}((x, y), \eta, \xi) &= h_2(x, y) |\xi| \xi, \end{aligned}$$

where $h(x, y) = 2e^{(x^2+y^2)}$. Let us consider the partial differential operator

$$Lu(x, y) = \Delta((x^2 + y^2)^{-1/2} |\Delta u| \Delta u) - \operatorname{div}((x^2 + y^2)^{-1/2} \mathcal{A}((x, y), u, \nabla u)).$$

Therefore, by Theorem 1, the problem

$$(P) \begin{cases} Lu(x) &= \frac{\cos(xy)}{(x^2 + y^2)} - \frac{\partial}{\partial x} \left(\frac{\sin(xy)}{(x^2 + y^2)} \right) - \frac{\partial}{\partial y} \left(\frac{\sin(xy)}{(x^2 + y^2)} \right), \quad \text{in } \Omega \\ u(x) &= 0, \quad \text{on } \partial\Omega \\ \Delta u &= 0, \quad \text{on } \partial\Omega \end{cases}$$

has a unique solution $u \in X = W^{2,3}(\Omega, \omega) \cap W_0^{1,3}(\Omega, \omega)$.

REFERENCES

- [1] A.C.Cavalheiro, *Existence and uniqueness of solutions for some degenerate nonlinear Dirichlet problems*, J. Appl. Anal., 19 (2013), 41-54, doi: 10.1515/jaa-2013-0003.
- [2] A.C.Cavalheiro, *Existence results for Dirichlet problems with degenerate p-Laplacian*, Opuscula Math., 33, no 3 (2013), 439-453, doi: 10.7494/OpMath.2013.33.3.439.
- [3] M. Chipot, *Elliptic Equations: An Introductory Course*, Birkhäuser, Berlin (2009).
- [4] P. Drábek, A. Kufner and F. Nicolosi, *Quasilinear Elliptic Equations with Degenerations and Singularities*, Walter de Gruyter, Berlin (1997).
- [5] E. Fabes, C. Kenig, R. Serapioni, *The local regularity of solutions of degenerate elliptic equations*, Comm. Partial Differential Equations 7, 1982, 77-116, doi:10.1080/03605308208820218.
- [6] S. Fučík, O. John and A. Kufner, *Function Spaces*, Noordhoff International Publ., Leyden, (1977).
- [7] J. Garcia-Cuerva and J.L. Rubio de Francia, *Weighted Norm Inequalities and Related Topics*, North-Holland Mathematics Studies 116, 1985.
- [8] D.Gilbarg and N.S. Trudinger, *Elliptic Partial Equations of Second Order*, 2nd Ed., Springer, New York (1983).
- [9] J. Heinonen, T. Kilpeläinen and O. Martio, *Nonlinear Potential Theory of Degenerate Elliptic Equations*, Oxford Math. Monographs, Clarendon Press, 1993.

- [10] A. Kufner, *Weighted Sobolev Spaces*, John Wiley & Sons, 1985.
- [11] B. Muckenhoupt, *Weighted norm inequalities for the Hardy maximal function*, *Trans. Amer. Math. Soc.* 165 ,1972, 207-226, doi:10.2307/1995882.
- [12] M. Talbi and N. Tsouli, *On the spectrum of the weighted p -Biharmonic operator with weight*, *Mediterr. J. Math.*, 4 (2007), 73-86, doi: 10.1007/s00009-007-0104-3.
- [13] A. Torchinsky, *Real-Variable Methods in Harmonic Analysis*, Academic Press, San Diego, 1986.
- [14] B.O. Turesson, *Nonlinear Potential Theory and Weighted Sobolev Spaces*, *Lecture Notes in Math.*, vol. 1736, Springer-Verlag, 2000.
- [15] E. Zeidler, *Nonlinear Functional Analysis and its Applications*, vol.I, Springer-Verlag, 1990.
- [16] E. Zeidler, *Nonlinear Functional Analysis and its Applications*, vol.II/B, Springer-Verlag, 1990.

A.C. CAVALHEIRO

DEPARTMENT OF MATHEMATICS, STATE UNIVERSITY OF LONDRINA, LONDRINA-PR, BRAZIL

E-mail address: accava@gmail.com